

On convergence of the Flint Hills series

Max A. Alekseyev*

January 20, 2013

Abstract

It is not known whether the Flint Hills series $\sum_{n=1}^{\infty} \frac{1}{n^3 \cdot \sin(n)^2}$ converges. We show that this question is closely related to the irrationality measure of π , denoted $\mu(\pi)$. In particular, convergence of the Flint Hills series would imply $\mu(\pi) \leq 2.5$ which is much stronger than the best currently known upper bound $\mu(\pi) \leq 7.6063 \dots$

This result easily generalizes to series of the form $\sum_{n=1}^{\infty} \frac{1}{n^u \cdot |\sin(n)|^v}$ where $u, v > 0$. We use the currently known bound for $\mu(\pi)$ to derive conditions on u and v that guarantee convergence of such series.

1 Introduction

Pickover [8] defined the *Flint Hills series* as $\sum_{n=1}^{\infty} \frac{1}{n^3 \cdot \sin(n)^2}$ (named after Flint Hills, Kansas) and questioned whether it converges. It was noticed that behavior of the partial sums of this series is closely connected to the rational approximations to π . In this paper we give a formal description of this connection, proving that convergence of the Flint Hills series would imply an upper bound 2.5 for the irrationality measure of π which is much stronger than the best currently known bound 7.6063... obtained by Salikhov [10]. A rather slow progress in evaluating the irrationality measure of π over past decades [6, 7, 2, 3, 4, 5, 10] indicates the hardness of this problem and suggests that the question of the Flint Hills series' convergence would unlikely be resolved in the nearest future.

The *irrationality measure* $\mu(x)$ of a positive real number x is defined as the infimum of such m that the inequality

$$0 < \left| x - \frac{p}{q} \right| < \frac{1}{q^m}$$

holds only for a finite number of co-prime positive integers p and q . If no such m exists, then $\mu(x) = +\infty$ (in which case x is called *Liouville number*).

Informally speaking, the larger is $\mu(x)$, the better x is approximated by rational numbers. It is known that $\mu(x) = 1$ if x is a rational number; $\mu(x) = 2$ if x is irrational algebraic number (Roth's theorem [9] for which Roth was awarded the Fields Medal); and $\mu(x) \geq 2$

*Department of Computer Science and Engineering, University of South Carolina, Columbia, SC, U.S.A.
Email: maxal@cse.sc.edu

if x is a transcendental number. Proving that $\mu(x) > 1$ is a traditional way to establish irrationality of x , with the most remarkable example of the $\zeta(3)$ irrationality (where $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$ is the Riemann zeta function) proved by Apéry [1, 11].

2 Convergence of the Flint Hills series

Lemma 1. *For a real number x , we have*

$$|\sin(x)| \leq |x|.$$

Furthermore, if $|x| \leq \pi/2$ then

$$|\sin(x)| \geq \frac{2}{\pi} \cdot |x|.$$

Proof. The former bound follows from the integral estimate

$$|\sin(x)| = \left| \int_0^x \cos y \cdot dy \right| \leq \int_0^{|x|} |\cos y| \cdot dy \leq \int_0^{|x|} 1 \cdot dy = |x|,$$

To prove the latter bound, we notice that $|\sin(x)| = \sin(|x|)$ and without loss of generality assume that $0 \leq x \leq \pi/2$. Let $x_0 = \arccos(2/\pi)$ so that for $x \leq x_0$ we have $\cos(x) \geq 2/\pi$ and thus

$$\sin(x) = \int_0^x \cos(y) \cdot dy \geq \int_0^x \frac{2}{\pi} \cdot dy = \frac{2}{\pi} \cdot x,$$

while for $x \geq x_0$ we have $\cos(x) \leq 2/\pi$ and thus

$$\sin(x) = 1 - \int_x^{\pi/2} \cos(y) \cdot dy \geq 1 - \int_x^{\pi/2} \frac{2}{\pi} \cdot dy = 1 - \frac{2}{\pi} \cdot \left(\frac{\pi}{2} - x \right) = \frac{2}{\pi} \cdot x.$$

□

Theorem 2. *For positive real numbers u and v , $\frac{1}{n^u \cdot |\sin(n)|^v} = O\left(\frac{1}{n^{u - (\mu(\pi) - 1) \cdot v - \epsilon}}\right)$ for any $\epsilon > 0$. Furthermore,*

1. *If $\mu(\pi) < 1 + u/v$, the sequence $\frac{1}{n^u \cdot |\sin(n)|^v}$ converges (to zero);*
2. *If $\mu(\pi) > 1 + u/v$, the sequence $\frac{1}{n^u \cdot |\sin(n)|^v}$ diverges.*

Proof. Let $\epsilon > 0$ and $k = \mu(\pi) + \epsilon/v$. Then the inequality

$$\left| \pi - \frac{p}{q} \right| < \frac{1}{q^k} \tag{1}$$

holds only for a finite number of co-prime positive integers p and q .

For a positive integer n , let $m = \lfloor n/\pi \rfloor$ so that $|n/\pi - m| \leq 1/2$ and thus $|n - m \cdot \pi| \leq \pi/2$. Then by Lemma 1,

$$|\sin(n)| = |\sin(n - m \cdot \pi)| \geq \frac{2}{\pi} \cdot |n - m \cdot \pi| = \frac{2}{\pi} \cdot m \cdot \left| \frac{n}{m} - \pi \right|.$$

On the other hand, for large enough n and m , we have $|n/m - \pi| \geq 1/m^k$, implying that

$$|\sin(n)| \geq \frac{2}{\pi} \cdot m \cdot \left| \frac{n}{m} - \pi \right| \geq \frac{2}{\pi} \cdot \frac{1}{m^{k-1}} \geq c \cdot \frac{1}{n^{k-1}}$$

for some constant $c > 0$ depending only on k but not n (since n/m tends to π as n grows).

Therefore, for all large enough n , we have

$$\frac{1}{n^u \cdot |\sin(n)|^v} \leq \frac{1}{c^v \cdot n^{u-(k-1)v}} = O\left(\frac{1}{n^{u-(\mu(\pi)-1)v-\epsilon}}\right).$$

The statement 1 now follows easily. If $\mu(\pi) < 1 + u/v$, we take $\epsilon = v/2 \cdot (1 + u/v - \mu(\pi))$ to obtain

$$\frac{1}{n^u \cdot |\sin(n)|^v} = O\left(\frac{1}{n^{u-v(\mu(\pi)-1)-\epsilon}}\right) = O\left(\frac{1}{n^\epsilon}\right).$$

Now let us prove statement 2. If $\mu(\pi) > 1 + u/v$, then for $k = 1 + u/v$ the inequality (1) holds for infinitely many co-prime positive integers p and q . That is, there exists a sequence of rationals p_i/q_i such that $|p_i - \pi \cdot q_i| < \frac{1}{q_i^{k-1}}$. Then

$$|\sin(p_i)| = |\sin(p_i - q_i \cdot \pi)| \leq |p_i - q_i \cdot \pi| < \frac{1}{q_i^{k-1}} < C \cdot \frac{1}{p_i^{k-1}}$$

for some constant $C > 0$ depending only on k .

Therefore, for $n = p_i$ we have

$$\frac{1}{n^u \cdot |\sin(n)|^v} > C^v \cdot n^{v(k-1)-u} = C^v.$$

On the other hand, we have

$$|\sin(1 + p_i)| = |\sin(1 + p_i - q_i \cdot \pi)| \xrightarrow{i \rightarrow \infty} \sin(1)$$

and thus

$$\frac{1}{(1 + p_i)^u \cdot |\sin(1 + p_i)|^v} \xrightarrow{i \rightarrow \infty} 0.$$

We conclude that the sequence $\frac{1}{n^u \cdot |\sin(n)|^v}$ diverges, since it contains two subsequences one which is bounded from below by a positive constant, while the other tends to zero. $\square$

Corollary 3. For positive real numbers u and v ,

1. If the sequence $\frac{1}{n^u \cdot |\sin(n)|^v}$ converges, then $\mu(\pi) \leq 1 + u/v$;
2. If the sequence $\frac{1}{n^u \cdot |\sin(n)|^v}$ diverges, then $\mu(\pi) \geq 1 + u/v$.

Corollary 4. If the Flint Hills series $\sum_{n=1}^{\infty} \frac{1}{n^3 \cdot \sin(n)^2}$ converges, then $\mu(\pi) \leq 5/2$.

Proof. Convergence of $\sum_{n=1}^{\infty} \frac{1}{n^3 \cdot \sin(n)^2}$ implies that $\lim_{n \rightarrow \infty} \frac{1}{n^3 \cdot \sin(n)^2} = 0$ and thus by Corollary 3, $\mu(\pi) \leq 5/2$. $\square$

Theorem 5. For positive real numbers u and v , if $\mu(\pi) < 1 + (u-1)/v$, then $\sum_{n=1}^{\infty} \frac{1}{n^u \cdot |\sin(n)|^v}$ converges.

Proof. The inequality $\mu(\pi) < 1 + (u-1)/v$ implies that $u - v \cdot (\mu(\pi) - 1) > 1$. Then there exists $\epsilon > 0$ such that $w = u - v \cdot (\mu(\pi) - 1) - \epsilon > 1$. By Theorem 2, $\frac{1}{n^u \cdot |\sin(n)|^v} = O\left(\frac{1}{n^w}\right)$ further implying that

$$\sum_{n=1}^{\infty} \frac{1}{n^u \cdot |\sin(n)|^v} = O(\zeta(w)) = O(1).$$

$\square$

Corollary 6. For positive real numbers u and v , if $\sum_{n=1}^{\infty} \frac{1}{n^u \cdot |\sin(n)|^v}$ diverges, then $\mu(\pi) \geq 1 + (u-1)/v$.

Unfortunately, the divergence of the Flint Hills series would not imply any non-trivial result per Corollary 6.

3 Known bounds for $\mu(\pi)$ and their implications

Since π is a transcendental number, $\mu(\pi) \geq 2$. To the best of our knowledge, no better lower bound for $\mu(\pi)$ is currently known.

The upper bound for $\mu(\pi)$ has been improved over the past decades. Starting with the bound $\mu(\pi) \leq 30$ established by Mahler in 1953 [6], it was improved to $\mu(\pi) \leq 20$ by Mignotte in 1974 [7], and then to $\mu(\pi) \leq 19.8899944\dots$ by Chudnovsky in 1982 [2]. In 1990-1993 Hata in a series of papers [3, 4, 5] decreased the upper bound down to $\mu(\pi) \leq 8.016045\dots$. The best currently known upper bound $\mu(\pi) \leq 7.6063\dots$ was obtained in 2008 by Salikhov [10].

By Theorem 2, the Salikhov's bound implies that the sequence $\frac{1}{n^u \cdot |\sin(n)|^v}$ converges to zero as soon as $1 + u/v > 7.6063$, including in particular the pairs $(u, v) = (7, 1), (14, 2), (20, 3)$ etc. Correspondingly, Theorem 5 further implies that the series $\sum_{n=1}^{\infty} \frac{1}{n^u \cdot |\sin(n)|^v}$ converges for $(u, v) = (8, 1), (15, 2), (21, 3)$ etc.

References

- [1] R. Apéry. Irrationalité de $\zeta(2)$ et $\zeta(3)$. *Asterisque*, 61:11–13, 1979.
- [2] G. V. Chudnovsky. Hermite-pade approximations to exponential functions and elementary estimates of the measure of irrationality of π . *Lecture Notes in Math.*, 925:299–322, 1982.
- [3] M. Hata. Legendre type polynomials and irrationality measures. *J. Reine Angew. Math.*, 407:99–125, 1990.
- [4] M. Hata. A lower bound for rational approximations to π . *J. Number Theor.*, 43(1):51–67, 1993.
- [5] M. Hata. Rational approximations to π and some other numbers. *Acta Arith.*, 63(4):335–349, 1993.
- [6] K. Mahler. On the approximation of π . *Nederl. Akad. Wetensch. Proc. Ser. A*, 56:30–42, 1953.

- [7] M. Mignotte. Approximations rationnelles de π et quelques autres nombres. *Bull. Soc. Math. Fr., Suppl.*, 37:121–132, 1974.
- [8] C. A. Pickover. *The Mathematics of Oz: Mental Gymnastics from Beyond the Edge*, volume 2. Cambridge University Press, 2002. Chapter 25 “Flint Hills Series”.
- [9] K. F. Roth. Rational approximations to algebraic numbers. *Mathematika*, 2:1–20, 168, 1955.
- [10] V. Kh. Salikhov. On the Irrationality Measure of π . *Russ. Math. Surv*, 63(3):570–572, 2008.
- [11] A. van der Poorten. A proof that euler missed... *The Mathematical Intelligencer*, 1(4):195–203, 1979.